

AD-A074 718 CARNEGIE-MELLON UNIV PITTSBURGH PA MANAGEMENT SCIENC--ETC F/G 12/1  
CONVEX PROGRAMS AND THEIR CLOSURES.(U)

OCT 78 C E BLAIR, J BORWEIN, R G JEROSLOW

N00014-75-C-0621


UNCLASSIFIED MSRR-430

NL

| OF |  
AD  
A074718




END  
DATE  
FILMED  
10-79  
DDC




MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 074718



*Handwritten signature or initials.*



**Carnegie-Mellon University**

PITTSBURGH, PENNSYLVANIA 15213


**LEVEL**

*See 1473 in locker.*

**GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION**


WILLIAM LARIMER MELLON, FOUNDER

*This document has been approved for public release and sale; its distribution is unlimited.*



DDC FILE COPY

79 10 01 066



Management Sciences Research Report No. 430

CONVEX PROGRAMS AND THEIR CLOSURES

by

C. E. Blair<sup>1</sup>

J. Borwein<sup>2</sup>

and

R. G. Jeroslow<sup>3</sup>

October 1978

This document has been approved  
for public release and sale; its  
distribution is unlimited.

<sup>1</sup> University of Illinois, Urbana

<sup>2</sup> Dalhousie University

<sup>3</sup> Carnegie-Mellon University and Georgia Institute of Technology

This report was prepared as part of the activities of the Management Sciences Research Group, Carnegie-Mellon University, and has been supported by NSF grant GP 37510 X1, and under Contract N00014-75-C-0621 NR 047-048 with the U. S. Office of Naval Research. Reproduction in whole or in part is permitted for any purpose of the U. S. Government.

Management Sciences Research Group  
Graduate School of Industrial Administration  
Carnegie-Mellon University  
Pittsburgh, Pennsylvania 15213

# Abstract

We extend the limiting lagrangean equation

*limit as theta approaches 0 (theta)* *sum over h as elements of H of*

$$\lim_{\theta \rightarrow 0^+} \sup_{\Lambda} \inf_{x \in K} (f_0^m(x) + \theta(w_0 x + w_1) + \sum_{h \in H} \lambda_h f_h^m(x)) = v(P) ,$$

*sup over lambda* *inf over x as elements of K* *lambda h*

and the results on affine supports from which it was deduced, to a very general setting that subsumes the previous constraint qualifications.

A simple example shows the need for some constraint qualification.

## Key Words:

- (1) Nonlinear programming
- (2) Lagrangean
- (3) Convexity



# CONVEX PROGRAMS AND THEIR CLOSURES

by

C. E. Blair,<sup>1</sup> J. Borwein,<sup>2</sup> and R. G. Jeroslow<sup>3</sup>

For a convex function  $f: D \rightarrow \mathbb{R}$  ( $D \subseteq \mathbb{R}^n$ ,  $D$  convex) the closure  $cl(f): cl(D) \rightarrow \mathbb{R} \cup \{+\infty\}$  is defined by

$$(1) \quad cl(f)(y) = \sup\{h(y) \mid h \text{ linear affine, } h(x) \leq f(x) \text{ for all } x \in D\}$$

where  $cl(D)$  is the closure of the convex set  $D$ .

It is well known that: (i)  $cl(f)(x) \leq f(x)$  for all  $x \in D$ ; (ii)  $cl(f)$  is convex; (iii)  $cl(f)(x) = f(x)$  for all  $x \in \text{relint}(D)$ , where  $\text{relint}(D)$  denotes the relative interior of the convex set  $D$ .

For a convex optimization problem (with possibly infinitely many constraints)

$$(P) \quad \begin{array}{l} \inf f_0(x) \\ \text{subject to } f_h(x) \leq 0 \text{ for } h \in H \\ \text{and } x \in K \end{array}$$

with optimal value denoted  $v(P)$ , the closure is

$$(P') \quad \begin{array}{l} \inf cl(f_0)(x) \\ \text{subject to } cl(f_h)(x) \leq 0 \text{ for } h \in H \\ \text{and } x \in cl(K) \end{array}$$

<sup>1</sup>University of Illinois, Urbana.

<sup>2</sup>Dalhousie University.

<sup>3</sup>Carnegie-Mellon University and Georgia Institute of Technology. This research has been supported by National Science Foundation grant GP 37510 X1, and Contract N00014-75-C-0621 NRO47-048 with the U.S. Office of Naval Research.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
<b>A</b>	

with optimal value denoted  $v(P')$ .

We assume throughout that  $(P)$  is consistent.

Duffin [1] and Jeroslow [2] show that when  $(P)$  and  $(P')$  have the same optimal value, a "limiting Lagrangean" exists, in the sense that (using the homotopy form of the limiting Lagrangean of [2, equation (50)])

$$(2) \quad \lim_{\theta \rightarrow 0+} \sup_{\Lambda} \inf_{x \in K} \{f_0(x) + \theta(wx + w_1) + \sum_{h \in H} \lambda_h f_h(x)\} = v(P)$$

for  $w \in \mathbb{R}^n$  and  $w_1 \in \mathbb{R}$  suitably chosen, where  $\Lambda$  denotes that space of vectors  $(\lambda_h | h \in H)$  which are nonnegative and only finitely non-zero. Moreover, from the value equality  $v(P) = v(P')$  also follows "fine detail" from which (2) is deduced, as e.g., [2, Theorem 3] and [2, Corollary 3].

It is also established in [2] that  $v(P) = v(P')$  holds in many instances in which the usual constraint qualifications, such as the existence of Slater points, may fail to hold even for  $|H|$  finite. This is because the limiting lagrangean (2) is not related to issues of linear affine, or even rather more general, supports to the perturbation function of  $(P)$ . This aspect of the limiting lagrangean was already present in the first limiting lagrangean, due to R. J. Duffin [1].

The purpose of this note, is to extend the validity of the limiting Lagrangean (and Theorem 3 and Corollary 3 of [2]) to a rather broad setting that is associated with the ordinary lagrangean in the case of  $|H|$  finite. We show, in this setting, that the limiting lagrangean holds again under weaker hypotheses than the ordinary lagrangean, even for  $|H|$  finite; and our result also treats  $|H|$  infinite.

Let  $D_h$  be the domain of definition of  $f_h$ . [2] showed that  $v(P) = v(P')$  if  $\text{relint}(K) \subseteq \text{relint}(D_h)$  for all  $h \in \{0\} \cup H$  and there was an  $x_0 \in \text{relint}(K)$  such that  $f_h(x_0) \leq 0$  for all  $h \in H$ . These latter hypotheses were called (CQ) in [2].

In this note we show:

**THEOREM:** Let  $H'$  denote those indices  $h \in \{0\} \cup H$  such that  $f_h$  is not closed.

$v(P) = v(P')$  if there is an  $x_0$  satisfying this constraint qualification:

$$(3) \quad x_0 \in \text{relint}(K) \cap \bigcap_{h \in H'} \text{relint}(D_h) \text{ and } f_h(x_0) \leq 0 \text{ for } h \in H.$$

(The intersection over an empty set is defined to be  $\mathbb{R}^n$ ).

**PROOF:** If  $x$  is feasible for (P) it is also feasible for (P') because  $\text{cl}(f_h)(x) \leq f_h(x)$ . Since  $\text{cl}(f_0)(x) \leq f_0(x)$ ,  $v(P') \leq v(P)$ .

To show that  $v(P') \geq v(P)$ , let  $x$  be any feasible point of (P'). For  $0 < \lambda < 1$  if  $y = \lambda x + (1 - \lambda)x_0$ ,  $y \in K$  and  $y \in \text{relint}(D_h)$  for all  $h \in H'$ , by the Accessibility Lemma [5, 3.2.11]. By (iii),  $\text{cl}(f_h)(y) = f_h(y)$  for all  $h \in H'$ ; therefore  $\text{cl}(f_h)(y) = f_h(y)$  for all  $h \in H \cup \{0\}$ .

Since  $\text{cl}(f_h)(x_0) \leq f_h(x_0) \leq 0$  and  $\text{cl}(f_h)(x) \leq 0$  for  $h \in H$ , one easily shows (by considering  $f_h$  and  $\text{cl}(f_h)$  on  $[x, x_0]$ ) that  $f_h(y) = \text{cl}(f_h)(y) \leq 0$  for  $h \in H$ . So  $y$  is a feasible point for (P).

By semi-continuity  $\lim_{\lambda \rightarrow 1} \text{cl}(f_0)(y) \leq \text{cl}(f_0)(x)$ . But since  $\text{cl}(f_0)(y) = f_0(y)$  for all  $\lambda < 1$  this implies  $v(P) \leq \text{cl}(f_0)(x)$ .

Since  $x$  was arbitrary, this shows  $v(P) \leq v(P')$ . Hence  $v(P) = v(P')$ , as desired.

Q.E.D.



REMARK: The same proof shows that, if  $K$  is closed, one obtains  $v(P) = v(P')$  from:

$$(4) \quad x_0 \in K \cap \bigcap_{h \in H'} \text{relint}(D_h)$$

$$\text{and } f_h(x_0) \leq 0 \text{ for } h \in H.$$

In one of the constraint qualifications of [2], it is assumed that  $H' = \emptyset$  and  $K$  is closed, in which case (4) becomes that constraint qualification  $(CQ)'$ . Trivially, (4) implies also the constraint qualification (CQ) of [2].

COROLLARY: Suppose that  $(P)$  has at least two different feasible points, and that none of the sets  $K$  or  $D_h$  for  $h \in H'$  contains any line segments in  $K \setminus \text{relint}(K)$  or  $D_h \setminus \text{relint}(D_h)$  (where  $H'$  is as defined in the theorem).

$$\text{Then } v(P) = v(P').$$

PROOF: Let  $x_a \neq x_b$  both be feasible in  $(P)$ . Since  $x_a, x_b \in K$  and  $K$  contains no line segment in  $K \setminus \text{relint}(K)$ ,  $x_0 = (x_a + x_b)/2 \in \text{relint}(K)$ .

Similarly,  $x_0 \in \text{relint}(D_h)$  for  $h \in H'$ . Trivially,  $f_h(x_0) \leq 0$  for  $h \in H$ .

The result now follows from the theorem.

Q.E.D.

Some constraint qualification is needed to insure that  $v(P) = v(P')$ , even for  $|H|$  finite. For example, consider this instance of a convex program:

$$\begin{aligned}
 (5) \quad & \inf x_1 \\
 & \text{subject to} \quad x_2 \leq 0 \\
 & f_2(x_1, x_2) \leq 0 \\
 & (x_1, x_2) \in K
 \end{aligned}$$

where  $K = \{(x_1, x_2) \mid 0 \leq x_1 \leq 1 \text{ and } x_2 \geq 0\}$  and

$$(6) \quad f_2(x_1, x_2) = \begin{cases} 0 & , \quad 0 \leq x_1 \leq 1 \text{ and } x_2 > 0; \\ 1 - x_1 & , \quad 0 \leq x_1 \leq 1 \text{ and } x_2 = 0; \\ +\infty & , \quad \text{otherwise.} \end{cases}$$

Here  $v(P) = 1$  and  $v(P') = 0$ , since  $\text{cl}(f_2)(x_1, x_2) \equiv 0$  if  $0 \leq x_1 \leq 1$  and  $x_2 \geq 0$ . In this example,  $H' = \{2\}$ , as  $f_2$  is not continuous on line segments that begin in the interior of  $K$  and end in the boundary segment  $\{(x_1, x_2) \mid x_2 = 0 \text{ and } 0 \leq x_1 < 1\}$ . Here also  $x_2 \leq 0$  and  $(x_1, x_2) \in K$  implies  $x_2 = 0$ , so  $(x_1, x_2) \notin \text{relint}(D_2)$ ; hence (3) fails.

August 2, 1978  
Revised September 14, 1978

## REFERENCES

- [1] R. J. Duffin, "Convex Analysis Treated by Linear Programming," Mathematical Programming 4 (1973), pp. 125-143.
- [2] R. G. Jeroslow, "A Limiting Lagrangean for Infinitely-Constrained Convex Optimization in  $R^n$ ," MSRR 417, GSIA, Carnegie-Mellon University, April 1978; revised June and July 1978.
- [3] O. L. Mangasarian, Nonlinear Programming, McGraw-Hill Book Co., New York, 1969.
- [4] R. T. Rockafellar, Convex Analysis, Princeton University Press, Princeton, New Jersey, 1970.
- [5] J. Stoer and C. Witzgall, Convexity and Optimization in Finite Dimensions: I, Springer Verlag, New York, 1970.



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 430	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>CONVEX PROGRAMS AND THEIR CLOSURES</b>	5. TYPE OF REPORT & PERIOD COVERED Technical Report October 1978	
6. AUTHOR(s) C. E. Blair, J. Borwein R. G. Jeroslow	7. PERFORMING ORG. REPORT NUMBER <b>MSRR-430</b>	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Graduate School of Industrial Administration Carnegie-Mellon University Pittsburgh, Pennsylvania 15213	9. CONTRACT OR GRANT NUMBER(s) <b>N00014-75-C-0621</b> <b>NSF-GP37548</b>	
10. CONTROLLING OFFICE NAME AND ADDRESS	11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 047-048	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. REPORT DATE <b>Oct 1978</b>	
	14. NUMBER OF PAGES 6	
	15. SECURITY CLASS. (of this report)	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  <b>9</b> Management sciences research rept.,		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Nonlinear programming, Lagrangean, Convexity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We extend the limiting Lagrangean equation $\lim_{\theta \rightarrow 0^+} \sup_{\Lambda} \inf_{x \in K} \{f_0(x) + \theta(wx + w_1) + \sum_{h \in H} \lambda_h f_h(x)\} = v(P),$ and the results on affine supports from which it was deduced, to a very general setting that subsumes the previous constraint qualifications.  A simple example shows the need for some constraint qualification.		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6001

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

403 426